# ACS Chemical Neuroscience

# Rethinking 5-HT<sub>1A</sub> Receptors: Emerging Modes of Inhibitory Feedback of Relevance to Emotion-Related Behavior

Stefanie C. Altieri,<sup>†,⊥</sup> Alvaro L. Garcia-Garcia,<sup>||,⊥</sup> E. David Leonardo,<sup>\*,||</sup> and Anne M. Andrews<sup>\*,†,‡,§</sup>

<sup>†</sup>Semel Institute for Neuroscience & Human Behavior, Hatos Center for Neuropharmacology, David Geffen School of Medicine, <sup>‡</sup>Department of Chemistry and Biochemistry, and <sup>§</sup>California NanoSystems Institute, University of California, Los Angeles, California 90095, United States

<sup>II</sup>Department of Psychiatry, Division of Integrative Neuroscience, Columbia University and New York State Psychiatric Institute, New York, New York 10032, United States

**Supporting Information** 

**ABSTRACT:** The complexities of the involvement of the serotonin transmitter system in numerous biological processes and psychiatric disorders is, to a substantial degree, attributable to the large number of serotonin receptor families and subtypes that have been identified and characterized for over four decades. Of these, the 5-HT<sub>1A</sub> receptor subtype, which was the first to be cloned and characterized, has received considerable attention based on its purported role in the etiology and treatment of mood and anxiety disorders. 5-HT<sub>1A</sub> receptors function both at presynaptic (autoreceptor) and postsynaptic (heteroreceptor) sites. Recent research has implicated distinct roles for these two populations of receptors in mediating emotion-related behavior. New concepts as to how 5-HT<sub>1A</sub> receptors function to control serotonergic tone throughout life



were highlights of the proceedings of the 2012 Serotonin Club Meeting in Montpellier, France. Here, we review recent findings and current perspectives on functional aspects of 5-HT<sub>1A</sub> auto- and heteroreceptors with particular regard to their involvement in altered anxiety and mood states.

KEYWORDS: Serotonin, autoreceptor, heteroreceptor, development, behavior, anxiety, antidepressant

# HISTORICAL PERSPECTIVE

Serotonin (5-hydroxytryptamine; 5-HT) is a monoamine neurotransmitter and neurohormone formed by the hydroxylation and subsequent decarboxylation of the essential dietary amino acid L-tryptophan. Serotonin is found primarily in the gastrointestinal tract, platelets, blood vessels, thyroid, pancreas, mammary glands,<sup>1–5</sup> and central nervous system (CNS).<sup>6,7</sup> In the brain, serotonin is thought to be a key modulatory neurotransmitter involved in the regulation of numerous physiological and behavioral processes including mood- and anxiety-related behavior, cognitive function, food intake, sexual behavior, sleep, cardiovascular function, blood pressure, pain, body temperature, and others.<sup>8–11</sup>

Serotonin was first reported in 1937 by Vialli and Erspamer and named enteramine.<sup>12,13</sup> In 1948, it was identified as a vasoconstrictor in blood serum where it was referred to as "serotonin".<sup>14</sup> Afterward, scientists realized that enteramine and serotonin were one in the same.<sup>15,16</sup> Serotonin was recognized as a neurotransmitter when it was discovered in extracts from mammalian brain.<sup>17,18</sup> In 1986, the pharmacology of serotonin was reviewed,<sup>19</sup> and for the first time the existence of three receptor families (5-HT1–3) was described; additional families were suspected. It is now known that the effects of serotonin are mediated by at least 14 different receptors, which are grouped into subfamilies based on pharmacological responses to specific ligands, sequence similarities at the gene and amino acid levels, gene organization, and second messenger coupling pathways.<sup>20–23</sup> Serotonin receptors are assigned to one of seven families, 5-HT<sub>1–7</sub>, with individual subtypes further designated by letters.

Among serotonin receptors, much attention has been focused on the 1A subtype  $(5-HT_{1A})$ . The human  $5-HT_{1A}$  receptor was cloned in 1987 as a single intronless gene<sup>24</sup> located on chromosome 5 (5q11.2-q13). In mice, the *Htr1a* gene resides on the distal part of chromosome 13. The  $5-HT_{1A}$  receptor protein consists of 422 amino acids. Evidence from human and rodent studies suggests that  $5-HT_{1A}$  receptors are implicated in a variety of physiological and pathological processes, such as learning, memory, schizophrenia, Parkinson's disease, and notably in the etiology and treatment of mood and anxiety disorders.<sup>25–32</sup>

Special Issue: Celebrating 25 Years of the Serotonin Club

Received:November 29, 2012Accepted:December 20, 2012Published:December 20, 2012





**Figure 1.** Inhibitory mechanisms of 5-HT<sub>1A</sub> autoreceptors. (A) Serotonergic cell bodies expressing 5-HT<sub>1A</sub> autoreceptors are located deep within the brainstem (blue box). Serotonin neurons projecting to the forebrain are organized into two main clusters designated as dorsal and median raphe nuclei with distinct subpopulations within these primary nuclei. 5-HT<sub>1A</sub> heteroreceptors are localized postsynaptically in various brain regions including the hippocampus (HIP), prefrontal cortex (PFC), thalamus (TH), lateral septum (SEP), amygdala (AMYG), and hypothalamic nuclei (HYP). Many of these regions have been associated with the pathophysiology of mood and anxiety disorders. (B) The top panel depicts a "conventional" serotonin synapse. In the bottom panel, activation of 5-HT<sub>1A</sub> autoreceptors controls serotonergic tone via one-to-one autoinhibitory feedback to reduce firing rates of serotonin neurons. 5-HT<sub>1A</sub> heteroreceptors also regulate serotonergic activity through descending glutamatergic projections originating in the medial prefrontal cortex (mPFC). This pathway makes connections with serotonin neurons via brainstem GABAergic inhibitory interneurons. (C) Current hypotheses regarding inhibitory mechanisms of 5-HT<sub>1A</sub> autoreceptor activation paint a more complex picture. For instance, emerging evidence suggests that serotonin neurons within a subpopulation or even across subpopulations affect each other via "lateral inhibition". Taken together, new perspectives on the functional aspects of 5-HT<sub>1A</sub> receptors associated with regulation of serotonergic activity are important avenues for future investigation, particularly regarding increased understanding of the roles of 5-HT<sub>1A</sub> receptors in the etiology and treatment of psychiatric disorders.

# ADULT EXPRESSION AND FUNCTION OF 5-HT<sub>1A</sub> RECEPTORS

Direct Autonomous Inhibition. In the mammalian brain, 5-HT<sub>1A</sub> receptors are divided into two distinct classes based on localization. The cell bodies of central serotonergic neurons are found in the raphe nuclei in the brain stem (Figure 1A),<sup>33,34</sup> where 5-HT<sub>1A</sub> receptors are located on soma and dendrites.<sup>35–37</sup> These 5-HT<sub>1A</sub> autoreceptors exert inhibitory feedback in response to local release of serotonin in the raphe nuclei from axonal collaterals (Figure 1B).38-43 Additionally, sources of local serotonin release have been proposed to arise from somatic and dendritic sites on serotonergic neurons.<sup>44–46</sup> Serotonin release in the cell body region results in reductions in serotonergic pacemaker activity and suppression of serotonin synthesis, turnover, and release in projection areas.<sup>47-51</sup> In support of this, constitutive 5-HT<sub>1A</sub> knockout mice show increased rates of serotonin neuronal discharge<sup>52</sup> and elevated basal dialysate serotonin levels in frontal cortex and hippocampus.53

Indirect Inhibition from Medial Prefrontal Cortex. 5-HT<sub>1A</sub> receptors are also expressed by nonserotonergic pyramidal, GABAergic, and cholinergic neurons<sup>54–58</sup> in limbic regions such as prefrontal cortex, hippocampus, lateral septum, and amygdala, as well as in several hypothalamic and thalamic

nuclei (Figure 1A).<sup>37,59,60</sup> Activation of 5-HT<sub>1A</sub> heteroreceptors mediates hyperpolarizing responses to released serotonin, which typically reduces postsynaptic neuronal excitability and firing rates.<sup>37,40,59</sup> As such, 5-HT<sub>1A</sub> heteroreceptors are involved in the modulation of other neurotransmission systems. For example, 5-HT<sub>1A</sub> receptors in the medial prefrontal cortex (mPFC) modulate dopamine cell firing and release.<sup>61–63</sup>

In addition to direct/local 5-HT<sub>1A</sub> receptor-mediated autoregulation, evidence exists for an indirect negative feedback mechanism that involves 5-HT<sub>1A</sub> heteroreceptors in the mPFC (Figure 1B).<sup>64–66</sup> The mPFC-dorsal raphe nucleus (DRN) pathway comprises glutamatergic descending projections that are hypothesized to decrease serotonin cell firing by activating DRN GABAergic interneurons to inhibit serotonin re-lease.<sup>65-72</sup> This pathway has been suggested as one of the neuroanatomical substrates controlling the effects of stress by preventing overactivation of DRN serotonergic neurons.<sup>73</sup> A minor glutamatergic pathway may also exist from the mPFC that excites serotonin cell firing directly in the DRN.<sup>68,69</sup> It is still not well understood how inhibitory 5-HT<sub>1A</sub> heteroreceptors excite cortical glutamatergic neurons. The most plausible explanation is the disinhibition of glutamatergic neurons via mPFC GABA interneurons expressing 5-HT<sub>1A</sub> heteroreceptors.<sup>58,74</sup> However, further studies are required to

determine the specific mechanisms of  $5\text{-HT}_{1A}$  receptors in the mPFC.

Intracellular Signaling. 5-HT<sub>1A</sub> receptors are inhibitory Gprotein coupled receptors (GPCRs). Early studies identified that 5-HT<sub>1A</sub> receptors function by coupling to Gi/Go proteins in most cells.<sup>75</sup> Extracellular receptor binding of serotonin or 5-HT<sub>1A</sub> agonists leads to intracellular exchange of GDP for GTP on Gi/Go alpha subunits. This, in turn, inhibits adenylyl cyclase, which reduces cAMP levels and protein kinase A activity.<sup>76</sup> Furthermore, agonist-induced activation of 5-HT1A receptors results in potassium channel activation and calcium channel inhibition.  $^{25,59,77-80}$  5-HT<sub>1A</sub> receptors are coupled to different GPCR pathways based on localization.<sup>81</sup> 5-HT<sub>1A</sub> autoreceptors couple exclusively to  $G\alpha_{i3}$  leading to partial inhibition of adenylyl cyclase,<sup>82–85</sup> an effect that may depend on specific agonists.<sup>83,86,87</sup> By contrast, 5-HT<sub>1A</sub> heteroreceptors couple mainly to  $G\alpha$ o in the hippocampus, and equally to  $G\alpha$ o and  $G\alpha$ i3 in cerebral cortex.<sup>88</sup> Differences in 5-HT<sub>1A</sub>  $G\alpha$ subunit coupling might explain regional differences in activation versus inhibition of intracellular signaling pathways. 5-HT<sub>1A</sub> autoreceptor desensitization is more pronounced compared to 5-HT<sub>1A</sub> heteroreceptor desensitization, which also could be related to differential G $\alpha$  coupling.<sup>88,89</sup> Stimulation of 5-HT<sub>1A</sub> receptors also leads to activation of G-protein-coupled inward rectifying potassium channels  $(GIRKs)^{90}$  in raphe neurons<sup>86,91–95</sup> and hippocampus.<sup>96–98</sup> Whether 5-HT<sub>1A</sub> receptors fully couple to inhibit adenylyl cyclase remains controversial, as it has been suggested that the hyperpolarizing response mediated by 5-HT<sub>1A</sub> autoreceptors is due to the activation of GIRK channels via G-protein  $\beta\gamma$  subunits.<sup>99</sup>

In addition to their canonical function, 5-HT<sub>1A</sub> receptors activate growth factor-regulated signaling pathways, such as mitogen-activated protein kinases (MAPK) and Akt signaling pathways.<sup>25</sup> In MAPK signaling, ERK is preferentially affected by 5-HT<sub>1A</sub> receptors. For example, in RN46A cells, a model of serotonergic raphe neurons that express endogenous 5-HT<sub>1A</sub> receptors, adenylyl cyclase and ERK1/2 phosphorylation were inhibited by 5-HT<sub>1A</sub>-receptor activation.<sup>100</sup> However, in hippocampal-derived differentiated HN2-5 cells, 5-HT<sub>1A</sub> agonists increased ERK phosphorylation and activity.<sup>101</sup> These and other studies suggest that the modulation of ERK may depend on neuronal origin, as well as maturation states.<sup>102–106</sup>

The Akt signaling pathway is also activated in 5-HT<sub>1A</sub> receptor-expressing cells and primary hippocampal neurons.<sup>104,107,108</sup> Activation of Akt by 5-HT<sub>1A</sub> receptors led to inactivation of GSK3 $\beta$  in hippocampal cultures,<sup>104,109,110</sup> an effect also observed in raphe cultures.<sup>105</sup> Recently, it has been suggested that the GSK3-regulating effects of 5-HT<sub>1A</sub> receptors are mediated by the PI3K/Akt signaling pathway.<sup>111</sup> Taken together, 5-HT<sub>1A</sub> autoreceptors and heteroreceptors have diverse intracellular signaling capabilities contributing to the complex regulation of the serotonin system, as well as neuronal networks modulated by serotonin.

### **5-HT<sub>1A</sub> RECEPTORS IN DEVELOPMENT**

Serotonin is a morphogenic factor,<sup>112</sup> and alteration of serotonin levels during early developmental windows has been shown to influence mood- and anxiety-related behavior in adult animals.<sup>113–115</sup> Thus, mapping and understanding factors that regulate the developmental trajectory of  $S-HT_{1A}$  receptor expression are important given that autoreceptors modulate serotonin release and heteroreceptors are widely

expressed in emotion-related circuits, where they mediate the effects of released serotonin. In rodents, 5-HT<sub>1A</sub> receptors appear in early embryonic development. Using in situ hybridization and immunocytochemistry, 5-HT<sub>1A</sub> mRNA has been detected as soon as embryonic day (E) 12 and 5-HT<sub>1A</sub> receptor protein by E14 in neuronal cultures prepared from brain stem.<sup>116</sup> A surge in 5-HT<sub>1A</sub> mRNA levels in the brain stem occurs beginning on E13, with peak levels occurring at E15-16.<sup>117</sup> Developmentally related changes in 5-HT<sub>1A</sub> receptor expression also occur at postsynaptic sites. Transient expression is evident in the septum and preoptic region during embryonic development.<sup>118</sup> 5-HT<sub>1A</sub> expression occurs at high levels in the cerebellum during the first two postnatal weeks tapering off to near undetectable levels by postnatal day (P) 21.119,120 In brain regions associated with the regulation of mood and anxiety, 5- $HT_{1A}$ -receptor expression likewise exhibits complex patterns during development. In situ hybridization and [<sup>3</sup>H]-8-OH-DPAT binding both indicate low 5-HT<sub>1A</sub> expression in the dentate gyrus granule cell layer of the hippocampus at E14.5, with levels gradually increasing during the first few postnatal weeks and reaching near adult levels by P13.<sup>118-121</sup>

Reports on the ontogeny of 5-HT<sub>1A</sub> receptors in the developing human brain, albeit fewer, are in agreement with those on rodents. In one study, [<sup>3</sup>H]-8-OH-DPAT binding was used to examine 5-HT<sub>1A</sub> receptor expression from tissue acquired at different stages of gestation.<sup>122</sup> Although expression levels varied with respect to gestational age (16-22 weeks), the relative regional distribution of 5-HT<sub>1A</sub> receptors was similar across time points. The highest receptor densities occurred in subregions of the hippocampus and frontal cortex, where a surge in 5-HT<sub>1A</sub> expression was observed at 18-22 weeks of gestation resulting in levels 3–4 times higher than those reported in adults.<sup>122</sup> Others have used in situ hybridization to compare 5-HT<sub>1A</sub> receptor mRNA levels in human tissue collected at  $\sim$ 28 weeks of gestation compared to 6–7 years of age.<sup>123</sup> Human fetal brain contained the highest 5-HT<sub>1A</sub> mRNA levels in raphe nuclei, cerebellum, and the CA1 and dentate gyrus regions of the hippocampus. When compared to later ages, only expression in the cerebellum showed dramatic changes, with mRNA levels being lower in children and undetectable in adults.<sup>123,124</sup> The latter is similar to the cerebellar expression pattern reported in rodents. Thus, 5-HT<sub>1A</sub>-receptor expression undergoes tight temporal regulation and in some regions, brief emergence and disappearance, suggesting morphogenic influences, especially in areas of the brain involved in modulating anxiety levels in adults.

A few studies have investigated functional aspects of  $5\text{-HT}_{1A}$  receptors during development by means of electrophysiology or pharmacologic challenge. Electrophysiological recordings in the prefrontal cortex of rats indicated that  $5\text{-HT}_{1A}$  receptor-mediated hyperpolarization appears late in postnatal development between P16 and P19 corresponding temporally with increases in receptor protein expression.<sup>118–120,125</sup> In mice,  $5\text{-HT}_{1A}$ -receptor mediated outward currents were not evident at P4 but were observed by P12.<sup>126</sup>  $5\text{-HT}_{1A}$  responses developed after the appearance of physiology characteristic of serotonergic neurons, which appeared to be fully developed by P12.  $5\text{-HT}_{1A}$  receptor-signaling pathways important for mediating developmental processes are complex. For example, 8-OH-DPAT-induced activation of the MAPK pathway, and specifically ERK1/2 kinases, requires different PKC isozymes to regulate distinct developmental processes (e.g., cell division

#### **ACS Chemical Neuroscience**

versus synaptic activity/strengthening) at P6 versus P15.<sup>127</sup> More work will be needed to understand functional changes in 5-HT<sub>1A</sub> mediated pre- and postsynaptic responses during postnatal development, particularly in terms of critical developmental windows important for shaping and determining anxiety-related behavior later in life.

# ■ 5-HT<sub>1A</sub> RECEPTORS IN MOOD AND ANXIETY

**Mechanisms of Antidepressants.** 5-HT<sub>1A</sub> autoreceptors play an important role in regulating serotonergic activity through feedback inhibition pathways. As such, the involvement of 5-HT<sub>1A</sub> receptors in the mechanism of action of antidepressants has been widely investigated, and literature supporting a role for this receptor population in the delayed efficacy of antidepressants has evolved. 5-HT<sub>1A</sub> autoreceptors (and possibly heteroreceptors) limit increases in extracellular serotonin levels induced by serotonin selective reuptake inhibitors (SSRIs). Even after chronic SSRI treatment, 5-HT<sub>1A</sub> autoreceptors maintain some control over serotonin release.<sup>128–132</sup>

The delayed efficacy of SSRIs has been partially attributed to the need for desensitization of 5-HT<sub>1A</sub> autoreceptors in the raphe nuclei, enabling firing rates of serotonergic neurons to overcome inhibition.<sup>128,133–137</sup> However, only ~50% of studies in rodents show increased extracellular serotonin after chronic administration of serotonin reuptake inhibiting antidepressants.<sup>138</sup> In this context, concomitant administration of 5-HT<sub>1A</sub> receptor antagonists with SSRIs has been hypothesized to hasten or potentiate changes in serotonin levels to improve clinical efficacy.<sup>54,139</sup> The ability of the partial 5-HT<sub>1A</sub> antagonist, pindolol, to accelerate antidepressant clinical efficacy has been investigated in patients with major depressive disorder. Meta-analyses suggest that pindolol administered in combination with SSRIs augments and accelerates symptomatic improvement after 2 weeks of treatment specifically in patients without prior history of treatment.<sup>140–142</sup>

However, not all findings support a role for 5-HT<sub>1A</sub> autoreceptor desensitization and its effects to accelerate SSRI treatment. For example, administration of pindolol in combination with SSRIs to patients with treatment resistant depression or extensive treatment histories does not hasten or produce symptom amelioration.<sup>143,144</sup> Human studies examining the role of 5-HT<sub>1A</sub> receptors in the treatment and etiology of major depressive disorder indicate that sensitivity to receptor signaling after administration of agonists are complicated based on reports of reduced receptor number and binding, as well as genetic influences.<sup>145</sup> Additionally, mechanisms of autoreceptor desensitization are not clear and may differ depending on the antidepressant administered. For example, chronic fluoxetine results in desensitization of autoreceptors, which is associated with a reduction in receptor stimulated [35S]GTPyS binding.<sup>146–149</sup> However, G-protein coupling is not altered in association with administration of the SSRI sertraline, indicating that antidepressants differ with regard to mechanisms of 5-HT<sub>1A</sub> desensitization.<sup>146</sup>

In addition to conflicting findings regarding 5-HT<sub>1A</sub> autoreceptor desensitization in clinical studies, desensitization occurs in animal models of depression generated via chronic stress paradigms.<sup>150–154</sup> 5-HT<sub>1A</sub> autoreceptors and heteroreceptors are also desensitized in serotonin transporter knockout mice in association with a phenotype characterized by enhanced anxiety.<sup>155–158</sup> Together, these studies highlight the complex regulation of 5-HT<sub>1A</sub> pathways, such that

desensitization may be common to both the treatment and precipitating etiological factors associated with major depressive disorder.

Animal Models of 5-HT<sub>1A</sub> Disruption. In light of data suggesting a role for 5-HT<sub>1A</sub> receptors in mediating antidepressant responses in humans, it is logical to suspect that these receptors also function in the neurocircuitry and pathophysiology of emotion-related behavior. Genetic manipulation of the murine Htr1A gene has been carried out to study the behavioral effects of receptor under- or overexpression. In both cases, the majority of data point to a role for  $5\text{-}\text{HT}_{1\text{A}}$ receptors in anxiety-related behavior. Table 1 in the Supporting Information provides a comprehensive summary of the literature to date focused on anxiety- and depression-like behaviors, and learning and memory in animal models of altered 5-HT<sub>1A</sub> expression. Taken together, these studies provide convincing evidence that global reductions in 5-HT<sub>1A</sub> receptor expression lead to increased anxiety-related behavior in adult animals across different strains of mice and behavior test paradigms.

The first reports on mice with reduced 5-HT<sub>1A</sub> expression appeared simultaneously in 1998. Three separate laboratories produced constitutive 5-HT<sub>1A</sub> knockout mice, each in a different background strain (Supporting Information Table 1).<sup>159–161</sup> Consistent changes in behavior were observed in tests for anxiety-related behavior associated with conflict in novel environments, including exploration in the center of the open field or the open arms of the elevated plus maze. Changes in behavior were more consistently replicated across tests in male versus female mice. In terms of depression-like behaviors, mice with reduced 5-HT<sub>1A</sub> expression displayed reduced immobility in the forced swim<sup>160,161</sup> and tail suspension tests.<sup>159</sup> However, interpretations of these behavioral changes were not in agreement. Some authors perceived decreased immobility as an increased stress response,<sup>160</sup> while others took a more traditional view of reduced immobility<sup>162</sup> as indicative of decreased learned helplessness/behavioral despair or an increased antidepressant-like response.<sup>159,160</sup>

Following these initial reports, others confirmed increased anxiety-like behavior in 5-HT<sub>1A</sub> knockout mice when presented with conflict situations (see Supporting Information Table 1 for a full list of references). Findings also included modestly enhanced fear responses to stressful stimuli such as foot shock.  $^{163-166}$  In terms of learning and memory, mice with reduced 5-HT<sub>1A</sub> expression displayed deficits in hippocampaldependent learning indicated by increased latencies and path lengths to find the hidden platform in the Morris water maze.<sup>167,168</sup> As opposed to mice with constitutive deletion of 5-HT<sub>1A</sub> receptors, mice with developmentally limited overexpression of 5-HT $_{1A}$  heteroreceptors in dentate gyrus and cortex also displayed reduced anxiety in novel, conflict environments.<sup>169</sup> By contrast, mice with permanent overexpression failed to show changes in anxiety-like behaviors.<sup>170</sup> Similar to  $5\text{-}HT_{1A}$  knockout mice, transient  $5\text{-}HT_{1A}$  overexpression resulted in deficits in hippocampal-dependent learning and memory in the Morris water maze.<sup>171</sup> Deficits were less pronounced or absent in permanent overexpressing mice.<sup>172</sup> Information pertaining to changes in learning and memory after 5-HT<sub>1A</sub> receptor manipulation is reviewed in detail elsewhere.<sup>172,173</sup>

Recently, studies have been carried out to delete specific subpopulations of 5-HT<sub>1A</sub> receptors enabling the differentiation of pre- versus postsynaptic behavioral effects.<sup>174</sup> When 5-HT<sub>1A</sub>

autoreceptors were selectively inactivated, mice exhibited increased anxiety-like behavior similar to constitutive knockout mice. These findings highlight the importance of raphe 5-HT<sub>1A</sub> receptors in mediating anxiety responses. By contrast, when 5-HT<sub>1A</sub> heteroreceptors were specifically targeted, normal anxiety-like behavior was observed; however, increased depression-like behavior was observed in the forced swim test.<sup>174</sup> Increased anhedonic behavior related to reduced consumption of palatable food has also been observed in mice with 5-HT<sub>1A</sub> heteroreceptor knockout. The latter findings on depression-related behavior illustrate an interesting dichotomy since mice with constitutive deletion of 5-HT<sub>1A</sub> auto- and heteroreceptors show decreases in depression-like behavior (see Supporting Information Table 1). Thus, the synergistic effects of genetically inactivating auto- and hetero 5-HT<sub>1A</sub> receptors cannot be predicted by studying knockout of either receptor population in isolation.

In addition to mice with complete loss of  $5\text{-HT}_{1A}$  autoreceptors, mice with 30% reductions in  $5\text{-HT}_{1A}$  autoreceptor expression have been generated.<sup>175</sup> These mice display potentiated responses to the administration of fluoxetine as indicated by increased dialysate serotonin levels. These findings are important in the context of human *HTR1A* gene polymorphisms, including the C(-1019)G *HTR1A* single nucleotide polymorphism,<sup>176-178</sup> which is hypothesized to affect the expression and function of 5-HT<sub>1A</sub> receptors and to contribute to the high numbers of antidepressant non-responders.<sup>179</sup>

# ■ NEW PERSPECTIVES ON 5-HT<sub>1A</sub> AUTORECEPTORS

Networks and 5-HT<sub>1A</sub> Inhibitory Function. It is widely accepted that 5-HT<sub>1A</sub> receptors located on serotonergic neurons participate in autoinhibition. However, a more complex picture of inhibitory function than simple one-toone regulation of serotonin cell firing is emerging. Serotonin neurons are anatomically heterogeneous and their projections to the forebrain are topographically organized.<sup>180-183</sup> Furthermore, distinct, reproducible patterns of serotonin neuron activation within the raphe nuclei can be seen in response to specific external stimuli, suggesting that anatomical topography underlies functional topography.<sup>182,184</sup> Additionally, microdialysis studies examining extracellular serotonin levels after the application of various stressors have supported the idea of defined network patterns of serotonin neurons such that stressinduced changes in serotonin levels are stressor- and region specific.<sup>185</sup> These observations suggest that feedback inhibitory pathways might work to control serotonin-network activity. In one model, raphe serotonin neurons exhibit autonomous feedback such that groups of functionally similar serotonergic neurons regulate themselves in a homeostatic manner (Figure 1C).<sup>180</sup>

Nonautonomous feedback has also been suggested in which crosstalk between distinct groups of serotonin neurons provides "lateral inhibition" influencing patterns of activation (Figure 1C).<sup>180</sup> Along these lines, recent work by Commons and Sperling has demonstrated complementary patterns of activation of serotonergic neurons after exposure to and withdrawal from nicotine, which are different and reciprocally switched after 5-HT<sub>1A</sub> receptor blockade.<sup>181</sup> These findings suggest that endogenous feedback inhibition provided by 5-HT<sub>1A</sub> receptors might be regionally organized and depend on behavioral states. While these observations do not exclude an autoregulatory function, they raise the possibility that

 $5-HT_{1A}$  receptors might also operate in a nonautonomous fashion to mediate communication between different groups of serotonin neurons, thus suggesting that  $5-HT_{1A}$  receptors are involved in a regional form of autoregulation between different raphe subfields. These studies are consistent with anatomical studies showing that there are many interconnections between the different raphe nuclei<sup>180</sup> and axon collaterals of serotonin neurons travel for some distance within the dorsal raphe nuclei.<sup>186</sup>

There are also a number of distinct serotonin subsystems with unique genetic programming and functions.<sup>187</sup> Recent studies have shown that, in raphe nuclei, there are at least three different serotonin cell-types grouped by anatomical, physiological, and molecular characteristics, and their distribution transcends the traditional anatomical classification of raphe subfields. Furthermore, forebrain regions receiving serotonergic projections are innervated by serotonin neurons with distinct characteristics, forming a highly organized circuit.<sup>188</sup> Therefore, identifying and understanding functionally specific axon collaterals within the raphe that are involved in controlling emotional behaviors is likely to be an important future direction of research.

# **FUTURE PROSPECTS**

**Targeting Specific Receptor Populations.** The evolving distinctions between the specific and diverse roles for 5-HT<sub>1A</sub> auto- versus heteroreceptors and the manner in which different populations of neurons expressing 5-HT<sub>1A</sub> receptors elicit control over serotonergic tone is now shifting away from simplified views centered on autonomous autoreceptor-mediated feedback inhibition. Classical definitions of auto-receptors, which have long been associated with controlling serotonergic firing rates via feedback inhibition, remain an important functional property of these receptors. Yet, rather than one-to-one associations, emerging findings point to the idea that inhibitory regulation is also more complex and might operate at several levels that involve multiple pathways and/or networks that influence neighboring circuits.

As the rich complexity of 5-HT<sub>1A</sub> receptor organization and function continues to be uncovered, optimizing treatment strategies for mood and anxiety disorders by preferentially targeting different 5-HT<sub>1A</sub> receptor populations or networks to produce therapeutic effects should become possible. For example, the novel 5-HT<sub>1A</sub> agonist, F15599, has been shown to reduce immobility in the forced swim test<sup>189</sup> and to promote these effects primarily through activation of 5-HT<sub>1A</sub> heteroreceptors localized in the frontal cortex.<sup>106</sup> These findings support the idea that in addition to 5-HT<sub>1A</sub> autoreceptor desensitization following antidepressant treatment, increased 5-HT<sub>1A</sub> heteroreceptor activity occurs to promote antidepressant efficacy.<sup>190–193</sup> Furthermore, in support of the roles of 5-HT<sub>1A</sub> autoreceptors in mood and anxiety-related behavior, recent advances using siRNA to silence 5-HT<sub>1A</sub> autoreceptor expression have been shown to ameliorate immobility associated with learned helplessness and to augment fluoxetine-induced increases in serotonin levels in postsynaptic regions.194,195

**5-HT<sub>1A</sub> Receptors: A View of the Future.** The serotonin system, and 5-HT<sub>1A</sub> receptors in particular, exhibit properties that continue to challenge and enlarge our understanding of receptor function. In addition to direct autonomous inhibition, evidence now exists for indirect inhibition of raphe serotonergic neurons arising from a mPFC-DRN pathway. It will be

#### **ACS Chemical Neuroscience**

informative to sort out whether direct and indirect pathways work independently or synergistically to limit serotonin release in the mPFC. Moreover, the actions of serotonin might be coordinated across different brain regions as part of a global circuitry via lateral inhibition within and across subpopulations of raphe neurons, some of which have different developmental origins and genetic lineages. Thus, a picture of 5-HT<sub>1A</sub> receptor-mediated control of emotion-related behavior via coordination of brain-wide networks is coming to light. Still, additional work will be needed to uncover precise interactions between different parts of this network, hierarchical principles of 5-HT<sub>1A</sub> receptor function, and early life sequencing with regard to lasting consequences for network development.

Many animal models of altered 5-HT<sub>1A</sub> receptor expression have been created. Taken together, studies on these models highlight two important aspects of 5-HT<sub>1A</sub> receptors in regard to anxiety-related behavior. First, most studies point to 5-HT<sub>1A</sub> receptors, particularly during postnatal development, as key regulators of anxiety (see Supporting Information Table 1). By contrast, pharmacologic inhibition or genetic manipulation of 5-HT<sub>1A</sub> receptors in adult animals fails to produce anxiogenic behavior.<sup>196,197</sup> Second, unraveling the significance of 5-HT<sub>1A</sub> auto- versus heteroreceptors in regard to establishing baseline anxiety behavior is important. Null 5-HT<sub>1A</sub> mutant mice in which 5-HT<sub>1A</sub> heteroreceptors were ectopically overexpressed during development<sup>169</sup> or where 5-HT<sub>1A</sub> receptor expression was restored in forebrain regions, for example, hippocampus, cortex, and striatum,<sup>196</sup> exhibit reversal or rescue of increased anxiety behavior, respectively. However, the importance of heteroreceptors in gain-of-function experiments has been questioned by recent tissue-specific conditional knockout strategies where 5-HT<sub>1A</sub> autoreceptors, but not heteroreceptors, have been shown to be critical components for establishing normal levels of anxiety.<sup>174</sup> As additional information on the specific roles of different 5-HT<sub>1A</sub> receptor subpopulations and their influence over emotion-related behaviors during various timeframes become clear, avenues for novel treatment strategies for mood and anxiety disorder should become evident.

Further, as we sort out which 5-HT<sub>1A</sub> receptor inhibitory circuits play key roles in shaping specific types of behavior, these are also expected to become targets for the development of more selective and, hopefully, more effective therapeutics. Recent findings have shown that different populations of 5-HT<sub>1A</sub> receptors are coupled to different/multiple intracellular signaling pathways. This suggests the possibility of developing allosteric modulators or other types of small-molecule drugs that modify specific intracellular signaling pathways in cases where a particular type of receptor is coupled to more than one pathway in the same cell type or to different  $G\alpha$  subunits in different cells types.

Finally, the serotonin system remains unique among known neurotransmitter systems in that it is the only system identified as having two molecularly distinct autoreceptors. In addition to 5-HT<sub>1A</sub> receptors, serotonergic function is regulated by 5-HT<sub>1B</sub> autoreceptors, which are expressed presynaptically on serotonin axon terminals, in addition to postsynaptic axonal localization associated with dopaminergic, GABAergic, and glutamatergic systems.<sup>198</sup> Thus, both 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptors function as autoreceptors and heteroreceptors. It is to think that 5-HT<sub>1B</sub> receptors might exhibit complexity similar to 5-HT<sub>1A</sub> receptors in terms of multiple inhibitory circuits or hierarchical organization. Perhaps, continued investigation of

5-HT<sub>1A</sub> (and 5-HT<sub>1B</sub>) receptors, as well as other receptor subtypes that function both as auto- and heteroreceptors, for example, dopamine D2 and noradrenergic alpha2 receptors, will reveal additional modes of regulatory feedback and network organization. Nonetheless, it appears that recent advances in understanding 5-HT<sub>1A</sub> receptor function lead the way in terms of extending and expanding our thinking as to how behavioral circuits are organized and controlled by a specific receptor subtype. They also provide exciting new opportunities for drug discovery and development, particularly for the treatment of mood and anxiety disorders.

#### ASSOCIATED CONTENT

#### **S** Supporting Information

Table detailing behavior phenotypes in models of 5-HT<sub>1A</sub> receptor disruption or overexpression. This material is available free of charge via the Internet at http://pubs.acs.org.

#### AUTHOR INFORMATION

#### **Corresponding Author**

\*(E.D.L.) Mailing address: Department of Psychiatry, 1051 Riverside Dr. Box 87, Columbia University, New York, NY 10032. E-mail: el367@columbia.edu. Phone: 212-543-5266. (A.M.A.) Neuroscience Research Building, Room 545, 635 Charles E. Young Dr. S, Box 957332, Los Angeles, CA 90095-7332. E-mail: aandrews@mednet.ucla.edu. Phone: 310-794-9421.

## **Author Contributions**

<sup>⊥</sup>These authors contributed equally.

#### Notes

The authors declare no competing financial interest.

#### REFERENCES

(1) Cote, F., Fligny, C., Bayard, E., Launay, J. M., Gershon, M. D., Mallet, J., and Vodjdani, G. (2007) Maternal serotonin is crucial for murine embryonic development. *Proc. Nat.l Acad. Sci. U.S.A. 104*, 329–334.

(2) Gershon, M. D. (2000) 5-HT (serotonin) physiology and related drugs. *Curr. Opin. Gastroenterol.* 16, 113–120.

(3) Paulmann, N., Grohmann, M., Voigt, J. P., Bert, B., Vowinckel, J., Bader, M., Skelin, M., Jevsek, M., Fink, H., Rupnik, M., and Walther, D. J. (2009) Intracellular serotonin modulates insulin secretion from pancreatic beta-cells by protein serotonylation. *PLoS Biol.* 7, e1000229.

(4) Walther, D. J., Peter, J. U., Bashammakh, S., Hortnagl, H., Voits, M., Fink, H., and Bader, M. (2003) Synthesis of serotonin by a second tryptophan hydroxylase isoform. *Science 299*, 76.

(5) Berger, M., Gray, J. A., and Roth, B. L. (2009) The expanded biology of serotonin. *Annu. Rev. Med.* 60, 355–366.

(6) Deneris, E. S., and Wyler, S. C. (2012) Serotonergic transcriptional networks and potential importance to mental health. *Nat. Neurosci.* 15, 519–527.

(7) Hensler, J. G. (2006) Serotonergic modulation of the limbic system. *Neurosci. Biobehav. Rev.* 30, 203–214.

(8) Jonnakuty, C., and Gragnoli, C. (2008) What do we know about serotonin? *J. Cell. Physiol.* 217, 301–306.

(9) Lam, D. D., Garfield, A. S., Marston, O. J., Shaw, J., and Heisler, L. K. (2010) Brain serotonin system in the coordination of food intake and body weight. *Pharmacol., Biochem. Behav.* 97, 84–91.

(10) Watts, S. W., Morrison, S. F., Davis, R. P., and Barman, S. M. (2012) Serotonin and blood pressure regulation. *Pharmacol. Rev.* 64, 359–388.

(11) Veasey, S. C., Fornal, C. A., Metzler, C. W., and Jacobs, B. L. (1995) Response of serotonergic caudal raphe neurons in relation to specific motor activities in freely moving cats. *J. Neurosci.* 15, 5346–5359.

(12) Vialli, M. E., V. (1937) Richerche sul secreto delle cellule enterocromaffini IX Intorno alla natura chimica della sostanza specifica. *Boll. Soc. Med.-Chir. Pavia 51*, 1111–1116.

(13) Vialli, M. E., V. (1940) Richerche di caratterizzazione chimica delle sostanze fenoliche presenti negli estratti acetonici di ghiandole salivari posteriori di Octopus vulgaris. *Arch. Fisiol.* 40, 293–302.

(14) Rapport, M. M., Green, A. A., and Page, I. H. (1948) Crystalline Serotonin. *Science* 108, 329–330.

(15) Erspamer, V., and Asero, B. (1952) Identification of enteramine, the specific hormone of the enterochromaffin cell system, as 5-hydroxytryptamine. *Nature 169*, 800–801.

(16) Erspamer, V., and Asero, B. (1953) Isolation of enteramine from extracts of posterior salivary glands of Octopus vulgaris and of Discoglossus pictus skin. *J. Biol. Chem.* 200, 311–318.

(17) Twarog, B. M., and Page, I. H. (1953) Serotonin content of some mammalian tissues and urine and a method for its determination. *Am. J. Physiol.* 175, 157–161.

(18) Amin, A. H., Crawford, T. B., and Gaddum, J. H. (1954) The distribution of substance P and 5-hydroxytryptamine in the central nervous system of the dog. *J. Physiol.* 126, 596–618.

(19) Bradley, P. B., Engel, G., Feniuk, W., Fozard, J. R., Humphrey, P. P., Middlemiss, D. N., Mylecharane, E. J., Richardson, B. P., and Saxena, P. R. (1986) Proposals for the classification and nomenclature of functional receptors for 5-hydroxytryptamine. *Neuropharmacology* 25, 563–576.

(20) Pytliak, M., Vargova, V., Mechirova, V., and Felsoci, M. (2011) Serotonin receptors - from molecular biology to clinical applications. *Physiol. Res.* 60, 15–25.

(21) Filip, M., and Bader, M. (2009) Overview on 5-HT receptors and their role in physiology and pathology of the central nervous system. *Pharmacol. Rep.* 61, 761–777.

(22) Hannon, J., and Hoyer, D. (2008) Molecular biology of 5-HT receptors. Behav. Brain Res. 195, 198–213.

(23) Nichols, D. E., and Nichols, C. D. (2008) Serotonin receptors. Chem. Rev. 108, 1614–1641.

(24) Kobilka, B. K., Frielle, T., Collins, S., Yang-Feng, T., Kobilka, T. S., Francke, U., Lefkowitz, R. J., and Caron, M. G. (1987) An intronless gene encoding a potential member of the family of receptors coupled to guanine nucleotide regulatory proteins. *Nature* 329, 75–79. (25) Polter, A. M., and Li, X. (2010) 5-HT1A receptor-regulated

signal transduction pathways in brain. *Cell. Signalling* 22, 1406–1412. (26) Akimova, E., Lanzenberger, R., and Kasper, S. (2009) The

serotonin-1A receptor in anxiety disorders. *Biol. Psychiatry 66,* 627–635.

(27) Gordon, J. A., and Hen, R. (2004) The serotonergic system and anxiety. *NeuroMol. Med. 5*, 27–40.

(28) Hirvonen, J., Karlsson, H., Kajander, J., Lepola, A., Markkula, J., Rasi-Hakala, H., Nagren, K., Salminen, J. K., and Hietala, J. (2008) Decreased brain serotonin 5-HT1A receptor availability in medicationnaive patients with major depressive disorder: An in-vivo imaging study using PET and [carbonyl-11C]WAY-100635. *Int. J. Neuropsychopharmacol.* 11, 465–476.

(29) Le Francois, B., Czesak, M., Steubl, D., and Albert, P. R. (2008) Transcriptional regulation at a HTR1A polymorphism associated with mental illness. *Neuropharmacology* 55, 977–985.

(30) Lesch, K. P., and Gutknecht, L. (2004) Focus on The 5-HT1A receptor: Emerging role of a gene regulatory variant in psychopathology and pharmacogenetics. *Int. J. Neuropsychopharmacol.* 7, 381–385.

(31) Ohno, Y. (2011) Therapeutic role of 5-HT1A receptors in the treatment of schizophrenia and Parkinson's disease. *CNS Neurosci. Ther.* 17, 58–65.

(32) Strobel, A., Gutknecht, L., Rothe, C., Reif, A., Mossner, R., Zeng, Y., Brocke, B., and Lesch, K. P. (2003) Allelic variation in 5-HT1A receptor expression is associated with anxiety- and depression-related personality traits. *J. Neural Transm. 110*, 1445–1453.

(33) Sotelo, C., Cholley, B., El Mestikawy, S., Gozlan, H., and Hamon, M. (1990) Direct Immunohistochemical Evidence of the Existence of 5-HT1A Autoreceptors on Serotoninergic Neurons in the Midbrain Raphe Nuclei. *Eur. J. Neurosci.* 2, 1144–1154.

(34) Jans, L. A., Riedel, W. J., Markus, C. R., and Blokland, A. (2007) Serotonergic vulnerability and depression: assumptions, experimental evidence and implications. *Mol. Psychiatry* 12, 522–543.

(35) Kia, H. K., Miquel, M. C., Brisorgueil, M. J., Daval, G., Riad, M., El Mestikawy, S., Hamon, M., and Verge, D. (1996) Immunocytochemical localization of serotonin1A receptors in the rat central nervous system. *J. Comp. Neurol.* 365, 289–305.

(36) Pazos, A., and Palacios, J. M. (1985) Quantitative autoradiographic mapping of serotonin receptors in the rat brain. I. Serotonin-1 receptors. *Brain Res.* 346, 205–230.

(37) Riad, M., Garcia, S., Watkins, K. C., Jodoin, N., Doucet, E., Langlois, X., el Mestikawy, S., Hamon, M., and Descarries, L. (2000) Somatodendritic localization of 5-HT1A and preterminal axonal localization of 5-HT1B serotonin receptors in adult rat brain. *J. Comp. Neurol.* 417, 181–194.

(38) Hjorth, S., and Sharp, T. (1991) Effect of the 5-HT1A receptor agonist 8-OH-DPAT on the release of 5-HT in dorsal and median raphe-innervated rat brain regions as measured by in vivo micro-dialysis. *Life Sci. 48*, 1779–1786.

(39) Meller, E., Goldstein, M., and Bohmaker, K. (1990) Receptor reserve for 5-hydroxytryptamine1A-mediated inhibition of serotonin synthesis: Possible relationship to anxiolytic properties of 5-hydroxytryptamine1A agonists. *Mol. Pharmacol.* 37, 231–237.

(40) Sprouse, J. S., and Aghajanian, G. K. (1986) (-)-Propranolol blocks the inhibition of serotonergic dorsal raphe cell firing by 5-HT1A selective agonists. *Eur. J. Pharmacol.* 128, 295–298.

(41) Verge, D., Daval, G., Patey, A., Gozlan, H., el Mestikawy, S., and Hamon, M. (1985) Presynaptic 5-HT autoreceptors on serotonergic cell bodies and/or dendrites but not terminals are of the 5-HT1A subtype. *Eur. J. Pharmacol.* 113, 463–464.

(42) Wang, R. Y., and Aghajanian, G. K. (1977) Inhibiton of neurons in the amygdala by dorsal raphe stimulation: Mediation through a direct serotonergic pathway. *Brain Res. 120*, 85–102.

(43) Pineyro, G., and Blier, P. (1999) Autoregulation of serotonin neurons: role in antidepressant drug action. *Pharmacol. Rev.* 51, 533–591.

(44) Colgan, L. A., Cavolo, S. L., Commons, K. G., and Levitan, E. S. (2012) Action potential-independent and pharmacologically unique vesicular serotonin release from dendrites. *J. Neurosci.* 32, 15737–15746.

(45) Colgan, L. A., Putzier, I., and Levitan, E. S. (2009) Activitydependent vesicular monoamine transporter-mediated depletion of the nucleus supports somatic release by serotonin neurons. *J. Neurosci.* 29, 15878–15887.

(46) de Kock, C. P., Cornelisse, L. N., Burnashev, N., Lodder, J. C., Timmerman, A. J., Couey, J. J., Mansvelder, H. D., and Brussaard, A. B. (2006) NMDA receptors trigger neurosecretion of 5-HT within dorsal raphe nucleus of the rat in the absence of action potential firing. *J. Physiol.* 577, 891–905.

(47) Jolas, T., Haj-Dahmane, S., Lanfumey, L., Fattaccini, C. M., Kidd, E. J., Adrien, J., Gozlan, H., Guardiola-Lemaitre, B., and Hamon, M. (1993) -)Tertatolol is a potent antagonist at pre- and postsynaptic serotonin S-HT1A receptors in the rat brain. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 347, 453–463.

(48) Bohmaker, K., Eison, A. S., Yocca, F. D., and Meller, E. (1993) Comparative effects of chronic 8-OH-DPAT, gepirone and ipsapirone treatment on the sensitivity of somatodendritic 5-HT1A autoreceptors. *Neuropharmacology* 32, 527–534.

(49) Kennett, G. A., Marcou, M., Dourish, C. T., and Curzon, G. (1987) Single administration of 5-HT1A agonists decreases 5-HT1A presynaptic, but not postsynaptic receptor-mediated responses: Relationship to antidepressant-like action. *Eur. J. Pharmacol.* 138, 53–60.

(50) Blier, P., and de Montigny, C. (1987) Modification of 5-HT neuron properties by sustained administration of the 5-HT1A agonist gepirone: Electrophysiological studies in the rat brain. *Synapse 1*, 470– 480.

(51) Sprouse, J. S., and Aghajanian, G. K. (1988) Responses of hippocampal pyramidal cells to putative serotonin 5-HT1A and

5-HT1B agonists: A comparative study with dorsal raphe neurons. *Neuropharmacology* 27, 707–715.

(52) Richer, M., Hen, R., and Blier, P. (2002) Modification of serotonin neuron properties in mice lacking 5-HT1A receptors. *Eur. J. Pharmacol.* 435, 195–203.

(53) Parsons, L. H., Kerr, T. M., and Tecott, L. H. (2001) 5-HT(1A) receptor mutant mice exhibit enhanced tonic, stress-induced and fluoxetine-induced serotonergic neurotransmission. *J. Neurochem.* 77, 607–617.

(54) Artigas, F., Adell, A., and Celada, P. (2006) Pindolol augmentation of antidepressant response. *Curr. Drug Targets* 7, 139–147.

(55) Azmitia, E. C., Gannon, P. J., Kheck, N. M., and Whitaker-Azmitia, P. M. (1996) Cellular localization of the 5-HT1A receptor in primate brain neurons and glial cells. *Neuropsychopharmacology* 14, 35–46.

(56) Cassel, J. C., and Jeltsch, H. (1995) Serotonergic modulation of cholinergic function in the central nervous system: Cognitive implications. *Neuroscience* 69, 1–41.

(57) Palchaudhuri, M., and Flugge, G. (2005) 5-HT1A receptor expression in pyramidal neurons of cortical and limbic brain regions. *Cell Tissue Res.* 321, 159–172.

(58) Santana, N., Bortolozzi, A., Serrats, J., Mengod, G., and Artigas, F. (2004) Expression of serotonin1A and serotonin2A receptors in pyramidal and GABAergic neurons of the rat prefrontal cortex. *Cereb. Cortex* 14, 1100–1109.

(59) Hamon, M., Gozlan, H., el Mestikawy, S., Emerit, M. B., Bolanos, F., and Schechter, L. (1990) The central 5-HT1A receptors: Pharmacological, biochemical, functional, and regulatory properties. *Ann. N.Y. Acad. Sci. 600*, 114–129 discussion 129–131..

(60) Beck, S. G., Choi, K. C., and List, T. J. (1992) Comparison of 5hydroxytryptamine1A-mediated hyperpolarization in CA1 and CA3 hippocampal pyramidal cells. *J. Pharmacol. Exp. Ther.* 263, 350–359.

(61) Diaz-Mataix, L., Artigas, F., and Celada, P. (2006) Activation of pyramidal cells in rat medial prefrontal cortex projecting to ventral tegmental area by a 5-HT1A receptor agonist. *Eur. Neuropsychopharmacol.* 16, 288–296.

(62) Diaz-Mataix, L., Scorza, M. C., Bortolozzi, A., Toth, M., Celada, P., and Artigas, F. (2005) Involvement of 5-HT1A receptors in prefrontal cortex in the modulation of dopaminergic activity: Role in atypical antipsychotic action. *J. Neurosci.* 25, 10831–10843.

(63) Sakaue, M., Somboonthum, P., Nishihara, B., Koyama, Y., Hashimoto, H., Baba, A., and Matsuda, T. (2000) Postsynaptic 5hydroxytryptamine(1A) receptor activation increases in vivo dopamine release in rat prefrontal cortex. *Br. J. Pharmacol.* 129, 1028–1034.

(64) Sharp, T., Boothman, L., Raley, J., and Queree, P. (2007) Important messages in the 'post': recent discoveries in 5-HT neurone feedback control. *Trends Pharmacol. Sci.* 28, 629–636.

(65) Celada, P., Puig, M. V., Casanovas, J. M., Guillazo, G., and Artigas, F. (2001) Control of dorsal raphe serotonergic neurons by the medial prefrontal cortex: Involvement of serotonin-1A, GABA(A), and glutamate receptors. *J. Neurosci.* 21, 9917–9929.

(66) Hajos, M., Hajos-Korcsok, E., and Sharp, T. (1999) Role of the medial prefrontal cortex in 5-HT1A receptor-induced inhibition of 5-HT neuronal activity in the rat. *Br. J. Pharmacol.* 126, 1741–1750.

(67) Casanovas, J. M., Vilaro, M. T., Mengod, G., and Artigas, F. (1999) Differential regulation of somatodendritic serotonin 5-HT1A receptors by 2-week treatments with the selective agonists alnespirone (S-20499) and 8-hydroxy-2-(Di-n-propylamino)tetralin: Microdialysis and autoradiographic studies in rat brain. J. Neurochem. 72, 262–272.

(68) Hajos, M., Richards, C. D., Szekely, A. D., and Sharp, T. (1998) An electrophysiological and neuroanatomical study of the medial prefrontal cortical projection to the midbrain raphe nuclei in the rat. *Neuroscience* 87, 95–108.

(69) Jankowski, M. P., and Sesack, S. R. (2004) Prefrontal cortical projections to the rat dorsal raphe nucleus: Ultrastructural features and associations with serotonin and gamma-aminobutyric acid neurons. *J. Comp. Neurol.* 468, 518–529.

(70) Stratz, T., Farber, L., Varga, B., Baumgartner, C., Haus, U., and Muller, W. (2001) Fibromyalgia treatment with intravenous tropisetron administration. *Drugs Exp. Clin. Res.* 27, 113–118.

(71) Varga, V., Szekely, A. D., Csillag, A., Sharp, T., and Hajos, M. (2001) Evidence for a role of GABA interneurones in the cortical modulation of midbrain 5-hydroxytryptamine neurones. *Neuroscience* 106, 783–792.

(72) Borsini, F., Ceci, A., Bietti, G., and Donetti, A. (1995) BIMT 17, a 5-HT1A receptor agonist/5-HT2A receptor antagonist, directly activates postsynaptic 5-HT inhibitory responses in the rat cerebral cortex. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 352, 283–290.

(73) Amat, J., Baratta, M. V., Paul, E., Bland, S. T., Watkins, L. R., and Maier, S. F. (2005) Medial prefrontal cortex determines how stressor controllability affects behavior and dorsal raphe nucleus. *Nat. Neurosci. 8*, 365–371.

(74) Aznar, S., Qian, Z., Shah, R., Rahbek, B., and Knudsen, G. M. (2003) The 5-HT1A serotonin receptor is located on calbindin- and parvalbumin-containing neurons in the rat brain. *Brain Res.* 959, 58–67.

(75) De Vivo, M., and Maayani, S. (1986) Characterization of the 5hydroxytryptamine1a receptor-mediated inhibition of forskolin-stimulated adenylate cyclase activity in guinea pig and rat hippocampal membranes. *J. Pharmacol. Exp. Ther.* 238, 248–253.

(76) Cooper, D. M., and Londos, C. (1982) GTP-dependent stimulation and inhibition of adenylate cyclase. *Horiz. Biochem. Biophys.* 6, 309–333.

(77) Albert, P. R., and Robillard, L. (2002) G protein specificity: Traffic direction required. *Cell. Signalling* 14, 407–418.

(78) Barnes, N. M., and Sharp, T. (1999) A review of central 5-HT receptors and their function. *Neuropharmacology* 38, 1083–1152.

(79) Haj-Dahmane, S., Hamon, M., and Lanfumey, L. (1991) K+ channel and 5-hydroxytryptamine1A autoreceptor interactions in the rat dorsal raphe nucleus: An in vitro electrophysiological study. *Neuroscience* 41, 495–505.

(80) Lanfumey, L., and Hamon, M. (2000) Central 5-HT(1A) receptors: Regional distribution and functional characteristics. *Nucl. Med. Biol.* 27, 429–435.

(81) Mannoury la Cour, C., El Mestikawy, S., Hanoun, N., Hamon, M., and Lanfumey, L. (2006) Regional differences in the coupling of 5hydroxytryptamine-1A receptors to G proteins in the rat brain. *Mol. Pharmacol.* 70, 1013–1021.

(82) Liu, Y. F., Ghahremani, M. H., Rasenick, M. M., Jakobs, K. H., and Albert, P. R. (1999) Stimulation of cAMP synthesis by Gi-coupled receptors upon ablation of distinct Galphai protein expression. Gi subtype specificity of the 5-HT1A receptor. *J. Biol. Chem.* 274, 16444–16450.

(83) Marazziti, D., Palego, L., Giromella, A., Mazzoni, M. R., Borsini, F., Mayer, N., Naccarato, A. G., Lucacchini, A., and Cassano, G. B. (2002) Region-dependent effects of flibanserin and buspirone on adenylyl cyclase activity in the human brain. *Int. J. Neuropsychopharmacol.* 5, 131–140.

(84) Palego, L., Giromella, A., Marazziti, D., Borsini, F., Naccarato, A. G., Giannaccini, G., Lucacchini, A., Cassano, G. B., and Mazzoni, M. R. (1999) Effects of postmortem delay on serotonin and (+)8-OH-DPAT-mediated inhibition of adenylyl cyclase activity in rat and human brain tissues. *Brain Res.* 816, 165–174.

(85) Valdizan, E. M., Castro, E., and Pazos, A. (2010) Agonistdependent modulation of G-protein coupling and transduction of 5-HT1A receptors in rat dorsal raphe nucleus. *Int. J. Neuropsychopharmacol.* 13, 835–843.

(86) Clarke, W. P., Yocca, F. D., and Maayani, S. (1996) Lack of 5hydroxytryptamine1A-mediated inhibition of adenylyl cyclase in dorsal raphe of male and female rats. *J. Pharmacol. Exp. Ther.* 277, 1259– 1266.

(87) Johnson, R. G., Fiorella, D., Winter, J. C., and Rabin, R. A. (1997) [3H]8-OH-DPAT labels a 5-HT site coupled to inhibition of phosphoinositide hydrolysis in the dorsal raphe. *Eur. J. Pharmacol.* 329, 99–106.

(88) Mannoury la Cour, C., Boni, C., Hanoun, N., Lesch, K. P., Hamon, M., and Lanfumey, L. (2001) Functional consequences of 5-HT transporter gene disruption on 5-HT(1a) receptor-mediated regulation of dorsal raphe and hippocampal cell activity. *J. Neurosci.* 21, 2178–2185.

(89) Chaput, Y., Blier, P., and de Montigny, C. (1986) In vivo electrophysiological evidence for the regulatory role of autoreceptors on serotonergic terminals. *J. Neurosci.* 6, 2796–2801.

(90) Luscher, C., Jan, L. Y., Stoffel, M., Malenka, R. C., and Nicoll, R. A. (1997) G protein-coupled inwardly rectifying K+ channels (GIRKs) mediate postsynaptic but not presynaptic transmitter actions in hippocampal neurons. *Neuron 19,* 687–695.

(91) Bayliss, D. A., Li, Y. W., and Talley, E. M. (1997) Effects of serotonin on caudal raphe neurons: Activation of an inwardly rectifying potassium conductance. *J. Neurophysiol.* 77, 1349–1361.

(92) Clarke, W. P., De Vivo, M., Beck, S. G., Maayani, S., and Goldfarb, J. (1987) Serotonin decreases population spike amplitude in hippocampal cells through a pertussis toxin substrate. *Brain Res.* 410, 357–361.

(93) Katayama, J., Yakushiji, T., and Akaike, N. (1997) Characterization of the K+ current mediated by 5-HT1A receptor in the acutely dissociated rat dorsal raphe neurons. *Brain Res.* 745, 283–292.

(94) Kelly, J. S., Larkman, P., Penington, N. J., Rainnie, D. G., McAllister-Williams, H., and Hodgkiss, J. (1991) Serotonin receptor heterogeneity and the role of potassium channels in neuronal excitability. *Adv. Exp. Med. Biol.* 287, 177–191.

(95) Loucif, A. J., Bonnavion, P., Macri, B., Golmard, J. L., Boni, C., Melfort, M., Leonard, G., Lesch, K. P., Adrien, J., and Jacquin, T. D. (2006) Gender-dependent regulation of G-protein-gated inwardly rectifying potassium current in dorsal raphe neurons in knock-out mice devoid of the S-hydroxytryptamine transporter. *J. Neurobiol.* 66, 1475– 1488.

(96) Andrade, R., and Nicoll, R. A. (1987) Pharmacologically distinct actions of serotonin on single pyramidal neurones of the rat hippocampus recorded in vitro. *J. Physiol.* 394, 99–124.

(97) Colino, A., and Halliwell, J. V. (1987) Differential modulation of three separate K-conductances in hippocampal CA1 neurons by serotonin. *Nature* 328, 73–77.

(98) Oleskevich, S. (1995) G alpha ol decapeptide modulates the hippocampal 5-HT1A potassium current. *J. Neurophysiol.* 74, 2189–2193.

(99) Kovoor, A., and Lester, H. A. (2002) Gi Irks GIRKs. *Neuron 33,* 6–8.

(100) Kushwaha, N., and Albert, P. R. (2005) Coupling of 5-HT1A autoreceptors to inhibition of mitogen-activated protein kinase activation via G beta gamma subunit signaling. *Eur. J. Neurosci.* 21, 721–732.

(101) Adayev, T., El-Sherif, Y., Barua, M., Penington, N. J., and Banerjee, P. (1999) Agonist stimulation of the serotonin1A receptor causes suppression of anoxia-induced apoptosis via mitogen-activated protein kinase in neuronal HN2–5 cells. *J. Neurochem.* 72, 1489–1496.

(102) Albert, P. R., and Tiberi, M. (2001) Receptor signaling and structure: Insights from serotonin-1 receptors. *Trends Endocrinol. Metab.* 12, 453–460.

(103) Chang, C. W., Poteet, E., Schetz, J. A., Gumus, Z. H., and Weinstein, H. (2009) Towards a quantitative representation of the cell signaling mechanisms of hallucinogens: Measurement and mathematical modeling of 5-HT1A and 5-HT2A receptor-mediated ERK1/2 activation. *Neuropharmacology* 56 (Suppl 1), 213–225.

(104) Cowen, D. S., Johnson-Farley, N. N., and Travkina, T. (2005) 5-HT receptors couple to activation of Akt, but not extracellularregulated kinase (ERK), in cultured hippocampal neurons. *J. Neurochem.* 93, 910–917.

(105) Druse, M., Tajuddin, N. F., Gillespie, R. A., and Le, P. (2005) Signaling pathways involved with serotonin1A agonist-mediated neuroprotection against ethanol-induced apoptosis of fetal rhombencephalic neurons. *Brain Res.* 159, 18–28.

(106) Newman-Tancredi, A., Martel, J. C., Assie, M. B., Buritova, J., Lauressergues, E., Cosi, C., Heusler, P., Bruins Slot, L., Colpaert, F. C., Vacher, B., and Cussac, D. (2009) Signal transduction and functional selectivity of F15599, a preferential post-synaptic 5-HT1A receptor agonist. *Br. J. Pharmacol.* 156, 338–353.

(107) Hsiung, S. C., Tamir, H., Franke, T. F., and Liu, K. P. (2005) Roles of extracellular signal-regulated kinase and Akt signaling in coordinating nuclear transcription factor-kappaB-dependent cell survival after serotonin 1A receptor activation. *J. Neurochem. 95*, 1653–1666.

(108) Hsiung, S. C., Tin, A., Tamir, H., Franke, T. F., and Liu, K. P. (2008) Inhibition of 5-HT1A receptor-dependent cell survival by cAMP/protein kinase A: Role of protein phosphatase 2A and Bax. *J. Neurosci. Res. 86*, 2326–2338.

(109) Chen, S., Owens, G. C., Crossin, K. L., and Edelman, D. B. (2007) Serotonin stimulates mitochondrial transport in hippocampal neurons. *Mol. Cell Neurosci.* 36, 472–483.

(110) Talbot, J. N., Jutkiewicz, E. M., Graves, S. M., Clemans, C. F., Nicol, M. R., Mortensen, R. M., Huang, X., Neubig, R. R., and Traynor, J. R. (2010) RGS inhibition at G(alpha)i2 selectively potentiates 5-HT1A-mediated antidepressant effects. *Proc. Natl. Acad. Sci. U.S.A.* 107, 11086–11091.

(111) Polter, A. M., Yang, S., Jope, R. S., and Li, X. (2012) Functional significance of glycogen synthase kinase-3 regulation by serotonin. *Cell. Signalling* 24, 265–271.

(112) Lauder, J. M., Wallace, J. A., Krebs, H., Petrusz, P., and McCarthy, K. (1982) In vivo and in vitro development of serotonergic neurons. *Brain Res. Bull. 9*, 605–625.

(113) Oberlander, T. F., Gingrich, J. A., and Ansorge, M. S. (2009) Sustained neurobehavioral effects of exposure to SSRI antidepressants during development: Molecular to clinical evidence. *Clin. Pharmacol. Ther.* 86, 672–677.

(114) Trowbridge, S., Narboux-Neme, N., and Gaspar, P. (2011) Genetic models of serotonin (5-HT) depletion: What do they tell us about the developmental role of 5-HT? *Anat. Rec.* 294, 1615–1623.

(115) van Kleef, E. S., Gaspar, P., and Bonnin, A. (2012) Insights into the complex influence of 5-HT signaling on thalamocortical axonal system development. *Eur. J. Neurosci.* 35, 1563–1572.

(116) Hillion, J., Catelon, J., Raid, M., Hamon, M., and De Vitry, F. (1994) Neuronal localization of 5-HT1A receptor mRNA and protein in rat embryonic brain stem cultures. *Brain. Res.* 79, 195–202.

(117) Hillion, J., Milne-Edwards, J. B., Catelon, J., de Vitry, F., Gros, F., and Hamon, M. (1993) Prenatal developmental expression of rat brain 5-HT1A receptor gene followed by PCR. *Biochem. Biophys. Res. Commun.* 191, 991–997.

(118) Bonnin, A., Peng, W., Hewlett, W., and Levitt, P. (2006) Expression mapping of 5-HT1 serotonin receptor subtypes during fetal and early postnatal mouse forebrain development. *Neuroscience 141*, 781–794.

(119) Daval, G., Verge, D., Becerril, A., Gozlan, H., Spampinato, U., and Hamon, M. (1987) Transient expression of 5-HT1A receptor binding sites in some areas of the rat CNS during postnatal development. *Int. J. Dev. Neurosci.* 5, 171–180.

(120) Miquel, M. C., Kia, H. K., Boni, C., Doucet, E., Daval, G., Matthiessen, L., Hamon, M., and Verge, D. (1994) Postnatal development and localization of 5-HT1A receptor mRNA in rat forebrain and cerebellum. *Brain Res. 80*, 149–157.

(121) Patel, T. D., and Zhou, F. C. (2005) Ontogeny of S-HT1A receptor expression in the developing hippocampus. *Brain Res.* 157, 42–57.

(122) Bar-Peled, O., Gross-Isseroff, R., Ben-Hur, H., Hoskins, I., Groner, Y., and Biegon, A. (1991) Fetal human brain exhibits a prenatal peak in the density of serotonin 5-HT1A receptors. *Neurosci. Lett.* 127, 173–176.

(123) del Olmo, E., Lopez-Gimenez, J. F., Vilaro, M. T., Mengod, G., Palacios, J. M., and Pazos, A. (1998) Early localization of mRNA coding for 5-HT1A receptors in human brain during development. *Brain Res.* 60, 123–126.

(124) del Olmo, E., Diaz, A., Guirao-Pineyro, M., del Arco, C., Pascual, J., and Pazos, A. (1994) Transient localization of 5-HT1A receptors in human cerebellum during development. Neurosci. Lett. 166, 149-152.

(125) Beique, J. C., Campbell, B., Perring, P., Hamblin, M. W., Walker, P., Mladenovic, L., and Andrade, R. (2004) Serotonergic regulation of membrane potential in developing rat prefrontal cortex: Coordinated expression of 5-hydroxytryptamine (5-HT)1A, 5-HT2A, and 5-HT7 receptors. *J. Neurosci.* 24, 4807–4817.

(126) Calizo, L. H., Piel, D. A., and Beck, S. G. (2010) Uncovering critical periods in the development of 5-HT neuron physiology and excitatory and inhibitory synaptic activity, Society for Neuroscience, San Diego, CA.

(127) Mehta, M., Ahmed, Z., Fernando, S. S., Cano-Sanchez, P., Adayev, T., Ziemnicka, D., Wieraszko, A., and Banerjee, P. (2007) Plasticity of 5-HT 1A receptor-mediated signaling during early postnatal brain development. *J. Neurochem.* 101, 918–928.

(128) Gardier, A. M., Malagie, I., Trillat, A. C., Jacquot, C., and Artigas, F. (1996) Role of 5-HT1A autoreceptors in the mechanism of action of serotoninergic antidepressant drugs: Recent findings from in vivo microdialysis studies. *Fundam. Clin. Pharmacol.* 10, 16–27.

(129) Hervas, I., Queiroz, C. M., Adell, A., and Artigas, F. (2000) Role of uptake inhibition and autoreceptor activation in the control of S-HT release in the frontal cortex and dorsal hippocampus of the rat. *Br. J. Pharmacol.* 130, 160–166.

(130) Hjorth, S., Westlin, D., and Bengtsson, H. J. (1997) WAY100635-induced augmentation of the 5-HT-elevating action of citalopram: Relative importance of the dose of the 5-HT1A (auto)receptor blocker versus that of the 5-HT reuptake inhibitor. *Neuropharmacology 36*, 461–465.

(131) Popa, D., Cerdan, J., Reperant, C., Guiard, B. P., Guilloux, J. P., David, D. J., and Gardier, A. M. (2010) A longitudinal study of 5-HT outflow during chronic fluoxetine treatment using a new technique of chronic microdialysis in a highly emotional mouse strain. *Eur. J. Pharmacol.* 628, 83–90.

(132) Trillat, A. C., Malagie, I., Mathe-Allainmat, M., Anmella, M. C., Jacquot, C., Langlois, M., and Gardier, A. M. (1998) Synergistic neurochemical and behavioral effects of fluoxetine and 5-HT1A receptor antagonists. *Eur. J. Pharmacol.* 357, 179–184.

(133) Artigas, F., Romero, L., de Montigny, C., and Blier, P. (1996) Acceleration of the effect of selected antidepressant drugs in major depression by 5-HT1A antagonists. *Trends Neurosci.* 19, 378–383.

(134) Blier, P., and de Montigny, C. (1983) Effects of quipazine on pre- and postsynaptic serotonin receptors: Single cell studies in the rat CNS. *Neuropharmacology* 22, 495–499.

(135) Blier, P., and De Montigny, C. (1983) Electrophysiological investigations on the effect of repeated zimelidine administration on serotonergic neurotransmission in the rat. J. Neurosci. 3, 1270–1278.

(136) Dawson, L. A., Nguyen, H. Q., Smith, D. I., and Schechter, L. E. (2000) Effects of chronic fluoxetine treatment in the presence and absence of  $(\pm)$ pindolol: A microdialysis study. *Br. J. Pharmacol.* 130, 797–804.

(137) El Mansari, M., Sanchez, C., Chouvet, G., Renaud, B., and Haddjeri, N. (2005) Effects of acute and long-term administration of escitalopram and citalopram on serotonin neurotransmission: An in vivo electrophysiological study in rat brain. *Neuropsychopharmacology* 30, 1269–1277.

(138) Luellen, B. A., Gilman, T. L., and Andrews, A. M. (2010) Presynaptic adaptive responses to constitutive versus adult pharmacologic inhibition of serotonin uptake. In *Experimental Models in Serotonin Transporter Research* (Kalueff, A. V. L., J.L., Ed.), Cambridge University Press, Cambridge.

(139) Artigas, F., Perez, V., and Alvarez, E. (1994) Pindolol induces a rapid improvement of depressed patients treated with serotonin reuptake inhibitors. *Arch. Gen. Psychiatry 51*, 248–251.

(140) Portella, M. J., de Diego-Adelino, J., Ballesteros, J., Puigdemont, D., Oller, S., Santos, B., Alvarez, E., Artigas, F., and Perez, V. (2011) Can we really accelerate and enhance the selective serotonin reuptake inhibitor antidepressant effect? A randomized clinical trial and a meta-analysis of pindolol in nonresistant depression. *J. Clin. Psychiatry* 72, 962–969. (141) Ballesteros, J., and Callado, L. F. (2004) Effectiveness of pindolol plus serotonin uptake inhibitors in depression: a metaanalysis of early and late outcomes from randomised controlled trials. *J. Affective Disord.* 79, 137–147.

(142) Whale, R., Terao, T., Cowen, P., Freemantle, N., and Geddes, J. (2010) Pindolol augmentation of serotonin reuptake inhibitors for the treatment of depressive disorder: a systematic review. *J. Psychopharmacol.* 24, 513–520.

(143) Portella, M. J., de Diego-Adelino, J., Puigdemont, D., Perez-Egea, R., Alvarez, E., Artigas, F., and Perez, V. (2009) Pindolol augmentation enhances response outcomes in first depressive episodes. *Eur. Neuropsychopharmacol.* 19, 516–519.

(144) Segrave, R., and Nathan, P. J. (2005) Pindolol augmentation of selective serotonin reuptake inhibitors: accounting for the variability of results of placebo-controlled double-blind studies in patients with major depression. *Hum. Psychopharmacol.* 20, 163–174.

(145) Savitz, J., Lucki, I., and Drevets, W. C. (2009) 5-HT(1A) receptor function in major depressive disorder. *Prog. Neurobiol.* 88, 17–31.

(146) Rossi, D. V., Burke, T. F., McCasland, M., and Hensler, J. G. (2008) Serotonin-1A receptor function in the dorsal raphe nucleus following chronic administration of the selective serotonin reuptake inhibitor sertraline. *J. Neurochem.* 105, 1091–1099.

(147) Pejchal, T., Foley, M. A., Kosofsky, B. E., and Waeber, C. (2002) Chronic fluoxetine treatment selectively uncouples raphe 5-HT(1A) receptors as measured by [(35)S]-GTP gamma S autoradiography. *Br. J. Pharmacol.* 135, 1115–1122.

(148) Shen, C., Li, H., and Meller, E. (2002) Repeated treatment with antidepressants differentially alters 5-HT1A agonist-stimulated [35S]GTP gamma S binding in rat brain regions. *Neuropharmacology* 42, 1031–1038.

(149) Castro, M., Diaz, A., del Olmo, E., and Pazos, A. (2003) Chronic fluoxetine induces opposite changes in G protein coupling at pre and postsynaptic 5-HT1A receptors in rat brain. *Neuropharmacology* 44, 93–101.

(150) Rozeske, R. R., Evans, A. K., Frank, M. G., Watkins, L. R., Lowry, C. A., and Maier, S. F. (2011) Uncontrollable, but not controllable, stress desensitizes 5-HT1A receptors in the dorsal raphe nucleus. *J. Neurosci.* 31, 14107–14115.

(151) Bambico, F. R., Nguyen, N. T., and Gobbi, G. (2009) Decline in serotonergic firing activity and desensitization of 5-HT1A autoreceptors after chronic unpredictable stress. *Eur. Neuropsychopharmacol.* 19, 215–228.

(152) Lanfumey, L., Pardon, M. C., Laaris, N., Joubert, C., Hanoun, N., Hamon, M., and Cohen-Salmon, C. (1999) 5-HT1A autoreceptor desensitization by chronic ultramild stress in mice. *NeuroReport 10*, 3369–3374.

(153) Grippo, A. J., Sullivan, N. R., Damjanoska, K. J., Crane, J. W., Carrasco, G. A., Shi, J., Chen, Z., Garcia, F., Muma, N. A., and Van de Kar, L. D. (2005) Chronic mild stress induces behavioral and physiological changes, and may alter serotonin 1A receptor function, in male and cycling female rats. *Psychopharmacology (Berlin, Ger.)* 179, 769–780.

(154) Evrard, A., Barden, N., Hamon, M., and Adrien, J. (2006) Glucocorticoid receptor-dependent desensitization of 5-HT1A autoreceptors by sleep deprivation: studies in GR-i transgenic mice. *Sleep* 29, 31–36.

(155) Holmes, A., Lit, Q., Murphy, D. L., Gold, E., and Crawley, J. N. (2003) Abnormal anxiety-related behavior in serotonin transporter null mutant mice: the influence of genetic background. *Genes Brain Behav.* 2, 365–380.

(156) Kalueff, A. V., Olivier, J. D., Nonkes, L. J., and Homberg, J. R. (2010) Conserved role for the serotonin transporter gene in rat and mouse neurobehavioral endophenotypes. *Neurosci. Biobehav. Rev.* 34, 373–386.

(157) Li, Q., Wichems, C., Heils, A., Lesch, K. P., and Murphy, D. L. (2000) Reduction in the density and expression, but not G-protein coupling, of serotonin receptors (5-HT1A) in 5-HT transporter knock-out mice: gender and brain region differences. J. Neurosci. 20, 7888–7895.

(158) Li, Q., Wichems, C., Heils, A., Van De Kar, L. D., Lesch, K. P., and Murphy, D. L. (1999) Reduction of 5-hydroxytryptamine (5-HT)(1A)-mediated temperature and neuroendocrine responses and 5-HT(1A) binding sites in 5-HT transporter knockout mice. *J. Pharmacol. Exp. Ther.* 291, 999–1007.

(159) Heisler, L. K., Chu, H. M., Brennan, T. J., Danao, J. A., Bajwa, P., Parsons, L. H., and Tecott, L. H. (1998) Elevated anxiety and antidepressant-like responses in serotonin 5-HT1A receptor mutant mice. *Proc. Natl. Acad. Sci. U.S.A.* 95, 15049–15054.

(160) Parks, C. L., Robinson, P. S., Sibille, E., Shenk, T., and Toth, M. (1998) Increased anxiety of mice lacking the serotonin1A receptor. *Proc. Natl. Acad. Sci. U.S.A.* 95, 10734–10739.

(161) Ramboz, S., Oosting, R., Amara, D. A., Kung, H. F., Blier, P., Mendelsohn, M., Mann, J. J., Brunner, D., and Hen, R. (1998) Serotonin receptor 1A knockout: An animal model of anxiety-related disorder. *Proc. Natl. Acad. Sci. U.S.A.* 95, 14476–14481.

(162) Porsolt, R. D., Bertin, A., Blavet, N., Deniel, M., and Jalfre, M. (1979) Immobility induced by forced swimming in rats: Effects of agents which modify central catecholamine and serotonin activity. *Eur. J. Pharmacol.* 57, 201–210.

(163) Groenink, L., van Bogaert, M. J., van der Gugten, J., Oosting, R. S., and Olivier, B. (2003) 5-HT1A receptor and 5-HT1B receptor knockout mice in stress and anxiety paradigms. *Behav. Pharmacol.* 14, 369–383.

(164) Gross, C., Santarelli, L., Brunner, D., Zhuang, X., and Hen, R. (2000) Altered fear circuits in 5-HT(1A) receptor KO mice. *Biol. Psychiatry* 48, 1157–1163.

(165) Klemenhagen, K. C., Gordon, J. A., David, D. J., Hen, R., and Gross, C. T. (2006) Increased fear response to contextual cues in mice lacking the 5-HT1A receptor. *Neuropsychopharmacology* 31, 101–111.

(166) Tsetsenis, T., Ma, X. H., Lo Iacono, L., Beck, S. G., and Gross, C. (2007) Suppression of conditioning to ambiguous cues by pharmacogenetic inhibition of the dentate gyrus. *Nat. Neurosci.* 10, 896–902.

(167) Sarnyai, Z., Sibille, E. L., Pavlides, C., Fenster, R. J., McEwen, B. S., and Toth, M. (2000) Impaired hippocampal-dependent learning and functional abnormalities in the hippocampus in mice lacking serotonin(1A) receptors. *Proc. Natl. Acad. Sci. U.S.A.* 97, 14731–14736.

(168) Wolff, M., Costet, P., Gross, C., Hen, R., Segu, L., and Buhot, M. C. (2004) Age-dependent effects of serotonin-1A receptor gene deletion in spatial learning abilities in mice. *Brain Res.* 130, 39–48.

(169) Kusserow, H., Davies, B., Hortnagl, H., Voigt, I., Stroh, T., Bert, B., Deng, D. R., Fink, H., Veh, R. W., and Theuring, F. (2004) Reduced anxiety-related behaviour in transgenic mice overexpressing serotonin 1A receptors. *Brain Res.* 129, 104–116.

(170) Bert, B., Fink, H., Hortnagl, H., Veh, R. W., Davies, B., Theuring, F., and Kusserow, H. (2006) Mice over-expressing the 5-HT(1A) receptor in cortex and dentate gyrus display exaggerated locomotor and hypothermic response to 8-OH-DPAT. *Behav. Brain Res.* 167, 328–341.

(171) Bert, B., Dere, E., Wilhelmi, N., Kusserow, H., Theuring, F., Huston, J. P., and Fink, H. (2005) Transient overexpression of the 5-HT1A receptor impairs water-maze but not hole-board performance. *Neurobiol. Learn. Mem.* 84, 57–68.

(172) Bert, B., Fink, H., Rothe, J., Walstab, J., and Bonisch, H. (2008) Learning and memory in 5-HT(1A)-receptor mutant mice. *Behav. Brain. Res.* 195, 78–85.

(173) Ogren, S. O., Eriksson, T. M., Elvander-Tottie, E., D'Addario, C., Ekstrom, J. C., Svenningsson, P., Meister, B., Kehr, J., and Stiedl, O. (2008) The role of 5-HT(1A) receptors in learning and memory. *Behav. Brain. Res.* 195, 54–77.

(174) Richardson-Jones, J. W., Craige, C. P., Nguyen, T. H., Kung, H. F., Gardier, A. M., Dranovsky, A., David, D. J., Guiard, B. P., Beck, S. G., Hen, R., and Leonardo, E. D. (2011) Serotonin-1A autoreceptors are necessary and sufficient for the normal formation of circuits underlying innate anxiety. *J. Neurosci.* 31, 6008–6018.

(175) Richardson-Jones, J. W., Craige, C. P., Guiard, B. P., Stephen, A., Metzger, K. L., Kung, H. F., Gardier, A. M., Dranovsky, A., David, D. J., Beck, S. G., Hen, R., and Leonardo, E. D. (2010) 5-HT1A autoreceptor levels determine vulnerability to stress and response to antidepressants. *Neuron* 65, 40–52.

(176) Blier, P. (2010) Altered function of the serotonin 1A autoreceptor and the antidepressant response. *Neuron* 65, 1–2.

(177) Lemonde, S., Du, L., Bakish, D., Hrdina, P., and Albert, P. R. (2004) Association of the C(-1019)G 5-HT1A functional promoter polymorphism with antidepressant response. Int J Neuropsychopharmacol 7, 501–506.

(178) Lemonde, S., Turecki, G., Bakish, D., Du, L., Hrdina, P. D., Bown, C. D., Sequeira, A., Kushwaha, N., Morris, S. J., Basak, A., Ou, X. M., and Albert, P. R. (2003) Impaired repression at a 5hydroxytryptamine 1A receptor gene polymorphism associated with major depression and suicide. *J. Neurosci.* 23, 8788–8799.

(179) Rush, A. J., Warden, D., Wisniewski, S. R., Fava, M., Trivedi, M. H., Gaynes, B. N., and Nierenberg, A. A. (2009) STAR\*D: Revising conventional wisdom. *CNS Drugs* 23, 627–647.

(180) Bang, S. J., Jensen, P., Dymecki, S. M., and Commons, K. G. (2012) Projections and interconnections of genetically defined serotonin neurons in mice. *Eur. J. Neurosci.* 35, 85–96.

(181) Sperling, R., and Commons, K. G. (2011) Shifting topographic activation and 5-HT1A receptor-mediated inhibition of dorsal raphe serotonin neurons produced by nicotine exposure and withdrawal. *Eur. J. Neurosci.* 33, 1866–1875.

(182) Waselus, M., Valentino, R. J., and Van Bockstaele, E. J. (2011) Collateralized dorsal raphe nucleus projections: a mechanism for the integration of diverse functions during stress. *J. Chem. Neuroanat.* 41, 266–280.

(183) Jensen, P., Farago, A. F., Awatramani, R. B., Scott, M. M., Deneris, E. S., and Dymecki, S. M. (2008) Redefining the serotonergic system by genetic lineage. *Nat. Neurosci.* 11, 417–419.

(184) Vasudeva, R. K., Lin, R. C., Simpson, K. L., and Waterhouse, B. D. (2011) Functional organization of the dorsal raphe efferent system with special consideration of nitrergic cell groups. *J. Chem. Neuroanat. 41*, 281–293.

(185) Adell, A., Casanovas, J. M., and Artigas, F. (1997) Comparative study in the rat of the actions of different types of stress on the release of 5-HT in raphe nuclei and forebrain areas. *Neuropharmacology 36*, 735–741.

(186) Li, Y. Q., Li, H., Kaneko, T., and Mizuno, N. (2001) Morphological features and electrophysiological properties of serotonergic and non-serotonergic projection neurons in the dorsal raphe nucleus. An intracellular recording and labeling study in rat brain slices. *Brain Res.* 900, 110–118.

(187) Gaspar, P., and Lillesaar, C. (2012) Probing the diversity of serotonin neurons. *Philos. Trans. R. Soc. London, Ser. B* 367, 2382–2394.

(188) Kiyasova, V., Bonnavion, P., Scotto-Lomassese, S., FabreV., Sahly, I., Toronche, F., Deneris, E., Gaspar, P., and Fernandez, S. P. (2012) A subpopulation of serotonergic neurons that do not express the 5-HT1A autoreceptor. *ACS Chem Neurosci.* DOI: 10.1021/cn300157s.

(189) Assie, M. B., Bardin, L., Auclair, A. L., Carilla-Durand, E., Depoortere, R., Koek, W., Kleven, M. S., Colpaert, F., Vacher, B., and Newman-Tancredi, A. (2010) F15599, a highly selective post-synaptic 5-HT(1A) receptor agonist: In-vivo profile in behavioural models of antidepressant and serotonergic activity. *Int. J. Neuropsychopharmacol.* 13, 1285–1298.

(190) Blier, P., Bergeron, R., and de Montigny, C. (1997) Selective activation of postsynaptic 5-HT1A receptors induces rapid antidepressant response. *Neuropsychopharmacology* 16, 333–338.

(191) Haddjeri, N., Blier, P., and de Montigny, C. (1998) Long-term antidepressant treatments result in a tonic activation of forebrain 5-HT1A receptors. J. Neurosci. 18, 10150–10156.

(192) Haddjeri, N., de Montigny, C., and Blier, P. (1999) Modulation of the firing activity of rat serotonin and noradrenaline neurons by  $(\pm)$ pindolol. *Biol. Psychiatry* 45, 1163–1169.

(193) Haddjeri, N., Ortemann, C., de Montigny, C., and Blier, P. (1999) Effect of sustained administration of the 5-HT1A receptor agonist flesinoxan on rat 5-HT neurotransmission. *Eur. Neuropsychopharmacol.* 9, 427–440.

(194) Bortolozzi, A., Castane, A., Semakova, J., Santana, N., Alvarado, G., Cortes, R., Ferres-Coy, A., Fernandez, G., Carmona, M. C., Toth, M., Perales, J. C., Montefeltro, A., and Artigas, F. (2012) Selective siRNA-mediated suppression of 5-HT1A autoreceptors evokes strong anti-depressant-like effects. *Mol Psychiatry* 17, 612–623.

(195) Ferres-Coy, A., Santana, N., Castane, A., Cortes, R., Carmona, M. C., Toth, M., Montefeltro, A., Artigas, F., and Bortolozzi, A. (2012) Acute 5-HT(1A) autoreceptor knockdown increases antidepressant responses and serotonin release in stressful conditions. *Psychopharmacology (Berlin, Ger.),*.

(196) Gross, C., Zhuang, X., Stark, K., Ramboz, S., Oosting, R., Kirby, L., Santarelli, L., Beck, S., and Hen, R. (2002) Serotonin1A receptor acts during development to establish normal anxiety-like behaviour in the adult. *Nature 416*, 396–400.

(197) Lo Iacono, L., and Gross, C. (2008) Alpha- $Ca^{2+}/calmodulin-$ dependent protein kinase II contributes to the developmental programming of anxiety in serotonin receptor 1A knock-out mice. *J. Neurosci.* 28, 6250–6257.

(198) Sari, Y. (2004) Serotonin1B receptors: from protein to physiological function and behavior. *Neurosci. Biobehav. Rev.* 28, 565–582.