

## Rostral brainstem contributes to medullary inhibition of muscle tone

J. M. SIEGEL, R. NIENHUIS and K. S. TOMASZEWSKI

*Neurobiology Research, V.A. Medical Center, Sepulveda, CA 91343 and Department of Psychiatry,  
UCLA School of Medicine, Los Angeles, CA 90024 (U.S.A.)*

(Accepted February 8th, 1983) *Key words:* medulla — pons — inhibition

— muscle tone — REM sleep

It has long been known that stimulation of the medial medulla in the decerebrate animal produces bilateral inhibition of muscle tone. In the present study we have found that transection of the brainstem at the ponto-medullary junction attenuates this inhibition. An interaction between medullary and rostral brainstem systems is responsible for the medullary inhibition phenomenon. A similar interaction may produce the inhibition of muscle tone seen in REM sleep.

Magoun and Rhines first demonstrated that stimulation of a large portion of the reticular formation of the medial medulla abolished tonic muscle activity and reflex response bilaterally in the response bilaterally in the decerebrate cat<sup>14</sup>. It was hypothesized that this effect was due to activation of an intrinsic medullary mechanism with projections descending to the spinal cord. A number of subsequent studies have shown that the medullary inhibition of muscle tone is mediated by hyperpolarization of motoneurons<sup>8,11,12</sup>.

A prolonged bilateral inhibition of muscle tone has been found to occur naturally in REM sleep<sup>9</sup>. This inhibition is also accompanied by hyperpolarization of motoneurons<sup>1-3,5,6,15,16,18</sup>. The motor inhibition of REM sleep can be prevented by small lesions placed in dorsolateral pons<sup>7,10,19</sup>. It has been hypothesized that this pontine region acts by projecting caudally to excite the medullary inhibitory region. Descending projections from the medullary region would then form the final common path for inhibition of motor activity.

If the medullary region responsible for atonia is autonomous, direct stimulation of this area, even after disconnection of the pons, should produce muscle atonia. However, we find that transection below the level of the pons greatly reduces the number of sites from which medullary inhibition can be produced. Interactions between the medulla and rostral brainstem contribute to the inhibition of muscle tone by medullary stimulation. These results have been reported in preliminary form<sup>20</sup>.

Eleven cats served as subjects. Eight were used in acute experiments. Four of the acute preparations were transected at the intercollicular level. The medulla was then stimulated to observe muscle tone inhibition. A second transection was performed at the ponto-medullary level in one of these cats and the stimulation procedures repeated. The 4 other acute preparations were transected only at the ponto-medullary level.

Three cats were used in chronic procedures. They were kept for 28 days after transection at the ponto-medullary level. The procedures for preparing and maintaining these animals are described elsewhere<sup>21</sup>. Two of these cats had 4 tripolar electrodes attached to implanted microdrives. These electrodes were used for stimulation between days 7 and 28 after the transection as they were advanced through the medullary inhibitory region. The other chronic cat was stimulated, following the same procedure used in the acute preparations (see below), in a terminal procedure on the 28th day after transection.

Surgical procedures prior to brainstem transection were carried out under halothane/oxygen anesthesia using aseptic technique. Midbrain transection was performed after removing the occipital lobe and hippocampus. A thin wedge of tissue was then aspirated in the coronal plane at the intercollicular or inferior collicular level.

Transection at the ponto-medullary junction was performed after aspirating the medial cerebellum to




Fig. 1. Effect of medullary stimulation on neck muscle tone recorded from electrodes placed in left (LFT) and right (RT) splenius muscles. Medullary stimulation produces suppression of muscle tone after decerebration and medial cerebellectomy. Transection at the pontomedullary junction prevents this suppression. Data are from a single experiment. Stimulation with 0.1 ms, 150  $\mu$ A pulses at 60 Hz for 300 ms was applied in each condition at P9.0, L1.0, H-8.0. Inhibition threshold was 80  $\mu$ A.

expose the floor of the fourth ventricle. A spatula was then lowered at 30° off vertical to transect the brainstem.

Blood pressure was monitored through the femoral artery after the more caudal transections, and remained above 100 mm Hg. Expired CO<sub>2</sub> was monitored in all cats through an endotracheal tube with a Beckman LB2 CO<sub>2</sub> analyzer, and respiration was assisted with a Harvard pump when necessary to maintain CO<sub>2</sub> levels below 6%. Core temperature was controlled by a circulating water heating pad triggered by a rectal thermistor. Pairs of stranded stainless steel wires, with 1 cm separation in the sagittal plane, were inserted into the left and right splenius muscles to monitor the activity of the dorsal neck musculature.

Tripolar 30-gauge stainless steel electrodes with 0.5 mm vertical tip separation were used for stimulation. Electrodes were lowered in 0.5 mm steps in the coronal plane. In acute preparations, stimulation was applied at  $L \pm 1.0$  and in the midline proceeding in a rostral to caudal direction. Stimulation was performed no less than 2 h after the cessation of halo-thane anesthesia. Medullary stimulation consisted of 300 or 500 ms trains of 0.1 ms pulses at 60 Hz. Stimulation intensity at each point was varied from 0 to 500  $\mu A$ . Current levels as high as 500  $\mu A$  were employed to rule out the possibility that transections had merely elevated the inhibition threshold. Muscle response was displayed on a Grass 78 polygraph and tape recorded along with blood pressure and percent CO<sub>2</sub>.

Medullary stimulation after intercollicular transection produced a complete bilateral suppression of muscle tone (Fig. 1), as first reported by Magoun<sup>13</sup>. Current thresholds were below 100  $\mu A$ . A rebound excitation was often evident immediately after stimulation. Aspiration of the medial cerebellum eliminated this rebound while the inhibitory effect remained (Fig. 1), as has previously been described<sup>22</sup>.

Transection at the ponto-medullary junction (Fig. 2), performed in both acute and chronic preparations, completely eliminated the bilateral suppression of muscle tone in 6 of the 8 transected cats. Stimulation of the region identified by Magoun and Rhines<sup>14</sup> produced only excitation of ipsilateral and contralateral neck musculature (Fig. 1) at current thresholds ranging from 80 to 160  $\mu A$ . Bilateral inhibition was never seen at any current level. In the re-




Fig. 2. Cat with midbrain and pontine transection. Electrode tracks in medulla pass through inhibitory area. After caudal transection, inhibition could not be evoked by medullary stimulation.

maintaining two cats, bilateral inhibition was found at less than 10% of the sites tested.

Histological reconstruction of the transection levels and stimulation sites was made for all brains. Transections at both midbrain and ponto-medullary junction were complete except for sparing of less than 0.5 mm of the lateral filaments of the branchium pontis in two cats. In the remaining cats a complete mechanical separation of the brainstem was effected at the transection sites. Results did not differ in these two groups. Ponto-medullary transection levels varied from the most caudal lesion passing through the abducens nucleus (P6.5, H-4) at a 30° angle to vertical, to the most rostral lesions passing at the same angle just caudal to the locus coeruleus complex (P5.4, H-2.8). The region tested for the inhibitory phenomenon extended from P7 to P16, H-5 to H-10, L.O.O  $\pm 1.0$ .

In their studies, Niemer and Magoun<sup>17</sup> reported that chronic hemisection of the brainstem at the ponto-medullary junction did not prevent the inhibition caused by stimulating either side of the medulla. They interpreted this finding as indicating that systems rostral to the medulla were not required for the inhibitory effect. The present results suggest a different interpretation. Rostral brainstem structures do contribute to medullary inhibition. However, unilateral connection of the pons to medullary structures is

- lary inhibition of muscle tone, *Soc. Neurosci. Abstr.*, 8 (1982)957.
- 21 Siegel, J. M., Nienhuis, R. and Tomaszewski, K. S., Behavioral states in the chronic medullary and mid-pontine cat, submitted.
- 22 Sprague, J. M. and Chambers, W. W., Control of posture by reticular formation and cerebellum in the intact, anesthetized and unanesthetized and in the decerebrated cat, *Amer. J. Physiol.*, 176 (1954) 52-64.
- 23 Tohyama, M., Sakai, K., Salvat, D., Touret, M. and Jouviet, M., Spinal projections from the lower brain stem in the cat as demonstrated by the horseradish peroxidase technique. I. Origins of the reticulo-spinal tracts and their funicular trajectories, *Brain Research*, 173 (1979)